CONGRESSO NAZIONALE SOCIETÀ ITALIANA FISSAZIONE ESTERNA

Fissazione esterna nel trattamento
delle emergenze e traumi militari,
tecniche di ricostruzione degli arti e
trattamento degli esiti posttraumatici

ROMA

2025

16-17 MAGGIO 2025

*Trattamento delle Osteomieliti in età pediatrica.
Ruolo della F.E.*

F.M. Pezzoli, A. Krzysztofiak*,
G. Mastantuoni, F. Falciglia

*UOC Ortopedia e Traumatologia
Dir. Francesco Falciglia*

**UOC Pediatria e Malattie Infettive
Resp. L.Lancella*

OSTEOMIELITI

Significativa **causa di morbilità** nei bambini in tutto il mondo (2-13/100.000 bambini/anno)

Possono avere alto tasso di **sequele** anche gravi e di lunga durata.

La **diagnosi tempestiva** ed un **trattamento adeguato** rimangono sempre di primaria importanza

Diagnosi differenziale con patologie tumorali !

Gruppo di studio multidisciplinare

**Infettivologi, Ortopedici, Radiologi,
Reumatologi, Microbiologi,
Chirurghi Plastici**

scientific reports

OPEN Clinical report and predictors of sequelae of 319 cases of pediatric bacterial osteomyelitis

Andrzej Krzysztofak^{1,11,12}, Marco Roversi^{2,3,11}, Antonio Musolino^{2,3}, Marco Cirillo⁴, Renato Maria Toniolo⁵, Osvaldo Mazza⁶, Livia Gargiullo⁷, Laura Lancella⁴, Paolo Rossi^{2,3}, Alberto Villani⁷ & Collaborative Osteomyelitis Study Group*

Pediatric osteomyelitis is an insidious disease that can lead to permanent sequelae, the management of which still relies on lengthy intravenous antibiotic therapy. The purpose of this study is to report and describe the clinical course and outcome of pediatric bacterial osteomyelitis in our experience. We reported the clinical, diagnostic, and treatment characteristics of all cases of osteomyelitis in children younger than 18 years of age who were hospitalized between January 2010 and December 2021 at the Bambino Gesù Children's Hospital in Rome, Italy, we compared patients with and without complications at follow-up, to identify any predictive factor for sequelae. The study sample included 319 cases of pediatric bacterial osteomyelitis. The median age was 7.77 years. Males (60.8%) were more affected than females. The most affected bones were the femur, tibia, and spine. Etiology was identified in 40.1% of cases, with *S.aureus* as the most common causative agent. Sequelae were reported in 43 cases (13.5%). The main predictors of sequelae were sepsis on admission and hypergammaglobulinemia. Our results show that a severe presentation with sepsis and hypergammaglobulinemia on admission may be associated with a higher frequency of late sequelae. Early recognition and aggressive treatment of this subgroup of patients may lead to a reduction in complications.

Gruppo di studio multidisciplinare

Casistica Osteomieliti OPBG 2010-2024

2022-2024 Gruppo di studio internazionale

COPYRIGHT © 2024 THE AUTHOR(S). PUBLISHED BY THE JOURNAL OF BONE AND JOINT SURGERY, INCORPORATED.

Prevalence of Complications Due to Transphyseal Hematogenous Osteomyelitis

Blaise Cochard, MD*, Asia Ciprani, MD*, Marco Cirillo, MD, Céline Habre, MD, Oscar Vazquez, MD, Louise Frizon, MD, Benedetta Bracci, MD, Romain Dayer, MD, Andrzej Krzysztofak, MD, and Dimitri Ceroni, MD

Background: Transphyseal hematogenous osteomyelitis (THO) is a common infectious condition, being present in 25% of patients with hematogenous osteomyelitis. A large proportion of pediatric hematogenous osteomyelitis infections can spread through the growth cartilage and therefore may be potentially responsible for growth disorders, leading to limb-length discrepancy or angular deformities. The purpose of the present study was to identify both the prevalence of complications caused by transphyseal osteomyelitis and factors influencing their occurrence.

Methods: The records for all patients who had been treated for THO over a 17-year period at the University Hospitals of Geneva and the Gesù Bambino Hospital in Rome were retrospectively analyzed. Clinical, biological, and bacteriological data were analyzed. Magnetic resonance imaging (MRI) scans were reviewed for all patients to assess the cross-sectional area of growth plate involvement. Restart of growth of the affected physeal cartilage was subsequently monitored by means of iterative radiographic examination.

Results: From a cohort of 594 patients with hematogenous osteomyelitis, 89 patients (15.0%) were found to have THO. The median age was 84 months (range, 1 to 167 months), with a bimodal distribution and peaks at 30 and 150 months; 59.6% (53) of the 89 patients were male. The lower limbs were most often affected, with the distal tibia and fibula accounting for 47.2% of all cases. Methicillin-sensitive *Staphylococcus aureus* and *Kingella kingae* were the most frequently identified pathogens, accounting for 61.8% of the cases. Virulence factors, such as Panton-Valentine leukocidin (PVL), toxic shock syndrome toxin (TSST), and accessory gene regulator (*Agr*), were recorded in 12.4% of MSSA strains. Transphyseal lesions affected a median of 7.6% (range, 0.79% to 58.2%) of the physeal surface. Complications affecting further growth were noted in 13% of patients with THO. Thrombocytopenia, leukocytosis, and the presence of virulence factors significantly influenced the occurrence of complications.

Conclusions: THO affects all age categories of the pediatric population. In the present study, growth disturbance occurred in 13% of cases. The presence of deep thrombocytopenia, leukocytosis, and virulence factors, such as *Agr*, TSST, and PVL, seems to strongly influence the occurrence of such complications.

Level of Evidence: Therapeutic Level IV. See Instructions for Authors for a complete description of levels of evidence.

Article

Multicenter Retrospective Analysis of 702 Pediatric Cases of Bone and Joint Infections: Definition of Clinical and Biological Features to Discriminate *K. kingae*, *S. aureus*, and Other Bacterial Infections

Marco Roversi ^{1,2}, Francesca Pignatelli ³, Giacomo De Marco ⁴ , Oscar Vazquez ⁴ , Dimitri Ceroni ⁴ , Antonio Musolino ⁵ , Marco Cirillo ⁶, Laura Lancella ³, Alberto Villani ^{5,7} , Andrzej Krzysztofak ^{3,*} and on behalf of the Osteomyelitis Collaborative Study Group [†]

¹ UOC Trials, Bambino Gesù Children's Hospital, IRCCS, 00165 Rome, Italy

² PhD Program in Immunology, Molecular Medicine and Applied Biotechnology, University of Rome Tor Vergata, 00133 Rome, Italy

³ Infectious Disease Unit, Bambino Gesù Children's Hospital, IRCCS, 00165 Rome, Italy; laura.lancella@opbg.net (L.L.)

⁴ Pediatric Orthopedic Unit, Pediatric Surgery Service, Geneva University Hospitals, 1206 Geneva, Switzerland

⁵ General Pediatrics and ED II Level, Bambino Gesù Children's Hospital IRCCS, 00165 Rome, Italy; antoniomusolino931@gmail.com (A.M.)

⁶ Diagnostic and Interventional Radiology Unit, Bambino Gesù Children's Hospital, IRCCS, 00165 Rome, Italy

⁷ System Medicine Department, Tor Vergata University, 00133 Rome, Italy

* Correspondence: andrzej.krzysztofak@opbg.net; Tel.: +39-066859-00165

† Collaborators of the Osteomyelitis Collaborative Study Group is provided in the Acknowledgments.

Casistica gruppo di studio multidisciplinare

Le localizzazioni

<i>Femore</i>	21.6%	79 pz
<i>Tibia</i>	15.8%	58 pz
<i>Colonna</i>	12.8%	47 pz

Linee Guida

Osteomieliti non complicate

 EUROPEAN SOCIETY for PAEDIATRIC INFECTIOUS DISEASES

BONE AND JOINT INFECTIONS

 Krzysztofak et al. *Italian Journal of Pediatrics* (2021) 47:179
<https://doi.org/10.1186/s13052-021-01130-4>

Italian Journal of Pediatrics

REVIEW

Open Access

Italian consensus on the therapeutic management of uncomplicated acute hematogenous osteomyelitis in children

Andrzej Krzysztofak^{1*}, Elena Chiappini², Elisabetta Venturini², Livia Gargiullo³, Marco Roversi¹, Carlotta Montagnani², Elena Bozzola³, Sara Chiurchiu¹, Davide Vecchio⁴, Elio Castagnola⁵, Paolo Tomà⁶, Gian Maria Rossolini⁷, Renato Maria Toniolo⁸, Susanna Esposito⁹, Marco Cirillo⁶, Fabio Cardinale¹⁰, Andrea Novelli¹¹, Giovanni Beltrami¹², Claudia Tagliabue¹³, Silvio Boero¹⁴, Daniele Deriu¹, Sonia Bianchini¹⁵, Annalisa Grandin³, Samantha Bosis¹³, Martina Ciarcia², Daniele Ciofi², Chiara Tersigni², Barbara Bortone², Giulia Trippella², Giangiacomo Nicolini¹⁶, Andrea Lo Vecchio¹⁷, Antonietta Giannattasio¹⁸, Paola Musso², Elena Serrano², Paola Marchisio¹³, Daniele Donà¹⁹, Silvia Garazzino²⁰, Luca Pierantoni²¹, Teresa Mazzone²², Paola Bernaschi²³, Alessandra Ferrari²⁴, Guido Castelli Gattinara²⁵, Luisa Galli² and Alberto Villani³

Journal of the Pediatric Infectious Diseases Society

GUIDELINES

Clinical Practice Guideline by the Pediatric Infectious Diseases Society and the Infectious Diseases Society of America: 2021 Guideline on Diagnosis and Management of Acute Hematogenous Osteomyelitis in Pediatrics

Charles R. Woods,¹ John S. Bradley,² Archana Chatterjee,³ Lawson A. Copley,⁴ Joan Robinson,⁵ Matthew P. Kronman,⁶ Antonio Arrieta,⁷ Sandra L. Fowler,⁸ Christopher Harrison,⁹ Maria A. Carrillo-Marquez,¹⁰ Sandra R. Arnold,¹¹ Stephen C. Eppes,¹¹ Laura P. Stadler,¹² Coburn H. Allen,¹³ Lynnette J. Mazur,¹⁴ C. Buddy Creech,¹⁵ Samir S. Shah,¹⁶ Theoklis Zaoutis,¹⁷ David S. Feldman,¹⁸ and Valéry Lavergne^{19,20}

Guideline Authors

Jesús Saavedra (Chair), Spain
 Oana Falup-Pecurariu, Romania
 Saul Faust, United Kingdom
 Hermann Girschick, Germany
 Nico Hartwig, Netherlands
 Sheldon Kaplan, United States
 Mathie Lorrot, France
 Elpis Mantadakis, Greece
 Heikki Peltola, Finland
 Pablo Rojo, Spain
 Theoklis Zaoutis, United States/Greece
 Anton LeMair, Netherlands

Importanza della diagnosi precoce

- La **diagnostica molecolare** deve essere utilizzata per identificare i batteri, utili per **un'appropriata terapia antibiotica**.
- È necessario un **approccio multidisciplinare**.

La **diagnostica molecolare** ha un ruolo sempre più importante nella diagnosi e gestione dell'osteomielite, offrendo strumenti per una identificazione più precisa dell'agente infettivo e per la valutazione dell'efficacia della terapia.

Pannello BIOFIRE® Joint Infection (JI)

Il pannello BIOFIRE® Joint Infection (JI) consente una rapida identificazione dei patogeni che causano comunemente infezioni articolari analizzando, in un unico test, 39 target inclusi batteri Gram-positivi, batteri Gram-negativi, lieviti e geni di resistenza antimicrobica.

- Semplice: 2 minuti di preparazione manuale
- Veloce: tempo di esecuzione di circa un'ora
- Completo: testa simultaneamente 39 patogeni e meccanismi di resistenza clinicamente rilevanti

Più veloce dei metodi tradizionali e più completo

INIZIO

Il Pannello BIOFIRE® JI ricerca 39 agenti patogeni e geni marcatori di resistenza antimicrobica che costituiscono la causa più frequente di infezioni articolari. Il test BIOFIRE® Joint Infection non è influenzato dal precedente uso di antibiotici e permette di rilevare anche batteri anaerobi di difficile crescita in coltura e organismi esigenti.

Rispetto ai metodi convenzionali, **migliora la resa diagnostica e facilita la rilevazione polimicrobica**.

Simultanea identificazione di 39 target

BATTERI Gram-Positivi	BATTERI Gram-Negativi
<i>Anaerococcus prevotii/vaginalis</i> <i>Clostridium perfringens</i> <i>Cutibacterium avidum/granulosum</i> <i>Enterococcus faecalis</i> <i>Enterococcus faecium</i> <i>Finegoldia magna</i> <i>Parvimonas micra</i> <i>Peptoniphilus</i> <i>Peptostreptococcus anaerobius</i> <i>Staphylococcus aureus</i> <i>Staphylococcus lugdunensis</i> <i>Streptococcus spp.</i> <i>Streptococcus agalactiae</i> <i>Streptococcus pneumoniae</i> <i>Streptococcus pyogenes</i>	<i>Bacteroides fragilis</i> <i>Citrobacter</i> <i>Enterobacter cloacae complex</i> <i>Escherichia coli</i> <i>Haemophilus influenzae</i> <i>Kingella kingae</i> <i>Klebsiella aerogenes</i> <i>Klebsiella pneumoniae group</i> <i>Morganella morganii</i> <i>Neisseria gonorrhoeae</i> <i>Proteus spp.</i> <i>Pseudomonas aeruginosa</i> <i>Salmonella spp.</i> <i>Serratia marcescens</i>
LIEVITI	GENI DI RESISTENZA Antimicrobica
<i>Candida spp.</i> <i>Candida albicans</i>	Carbapenemasi IMP KPC NDM OXA-48-like VIM ESBL CTX-M Resistenza alla Meticillina <i>mecA/C</i> and MREJ Resistenza alla Vancomicina <i>vanA/B</i>

Elimina le incertezze

INIZIO

I test convenzionali per le infezioni articolari sono complessi e spesso richiedono più di un singolo campione e giorni di attesa per l'ottenimento dei risultati. Il pannello BIOFIRE® Joint Infection è un test sindromico rapido che fornisce risultati completi disponibili in un report di facile interpretazione.

Utilizzando la tecnologia PCR, il pannello BIOFIRE® Joint Infection offre una risposta rapida ed accurata da singolo campione.

Osteomieliti complicate

Via ematica
Fratture esposte
Fratture conseguenti a traumi ad alta energia
Mezzi di sintesi

Casistica osteomieliti su fratture

N pat	Age years	sex	Days from trauma	Days recovery	pathogen	site	sequele	*K-wire
1	10	M	65	18	<i>Staph. aureus</i>	RADIUS ULNA	no	yes
2	15	M	4	15	ND	FEMUR	no	no
3	3	F	90	90	<i>Staph. Aureus</i> <i>MRSA</i>	TIBIA FIBULA	yes	yes
4	13	M	2	23	<i>Staph. aureus</i>	PELVIC GIRDLE	no	no
5	7	F	16	12	ND	CUNEIFORM	yes	no
6	17	M	13	50	<i>Enterobacter</i> <i>fecium</i>	FEMUR	yes	no
7	17	M	65	20	ND	ASTRAGALUS	no	no
8	2	M	32	23	<i>Staph. Aureus</i> <i>MRSA</i>	FEMUR	no	no
9	5	M	58	28	ND	TIBIA	no	no
10	12	M	7	25	<i>Klebsella p</i>	FEMUR	no	no
11	12	M	19	38	<i>Pseudomonas</i> <i>a.</i>	TIBIA	no	no
12	12	M	2	20	ND	FEMUR	no	no
13	9	F	20	67	<i>Staph.aureus</i>	RADIUS ULNA	no	yes
14	7	M	30	38	<i>Staph.aureus</i>	HUMERUS	yes	yes

Table 1. Clinical and Microbiological characteristics of patients with osteomyelitis after fracture. * K-wire =Kirschner wire

Osteomieliti post - traumatiche

E' un problema mondiale nel campo dei traumi ortopedici.

Non esiste un trattamento ideale per la sua gestione.

Questioni aperte:

- ✓ **il meccanismo fisiopatologico**
- ✓ **classificazione clinica, eziologica, istologica, anatomica**
- ✓ **diagnosi di imaging**
- ✓ **strategie terapeutiche**

Osteomieliti post - traumatiche

- rara nei bambini ma potenzialmente devastante con **esiti invalidanti**
- **La diagnosi** spesso difficile (anamnesi, clinica, microbiologia, dati di laboratorio, radiografia standard, TC, RM o scintigrafia ossea)
- Patogeni: *Staphylococcus Aureus* (incluso MRSA)

C. M. 7 aa - Frattura sovracondiloidea omero destro
Alterazione della struttura ossea diffusa del terzo diafisario e della metafisi distale dell'omero dx con apposizione periostale da osteomielite.

16-17 MAGGIO 2025

Ruolo della Fissazione Esterna

Stabilità
Non invasivo

Versatilità
Pulizia del focolaio

Terapia medica adeguata

16-17 MAGGIO 2025

Paziente 17 aa osteomielite da *Salmonella in Talassemia*

16-17 MAGGIO 2025

stabilita' meccanica

16-17 MAGGIO 2025

**Sequestrotomia
pulizia del focolaio,
pervieta' del canale**

ROMA

CONGRESSO NAZIONALE
SOCIETÀ ITALIANA
FISSAZIONE ESTERNA

16-17 MAGGIO 2025

**Invio del materiale asportato
per istologia e microbiologia**

Introduzione di preparato antibiotato

16-17 MAGGIO 2025

FUP

16-17 MAGGIO 2025

Pz 6 aa ricoverata per polmonite presenta successivamente tumefazione gamba sinistra

16-17 MAGGIO 2025

RX dopo un mese

16-17 MAGGIO 2025

Primo intervento chirurgico

**Peggioramento del quadro clinico
Trattamento non efficace**

secondo intervento montaggio fissatore esterno

16-17 MAGGIO 2025

Rx controlli successivi

16-17 MAGGIO 2025

RX dopo rimozione del fissatore esterno

prima

Osteomielite in trauma ad alta energia

Bambina di 4 aa infezione cute e tessuti molli associata a frattura pluri frammentaria scomposta della gamba e del piede.

16-17 MAGGIO 2025

Accorciamento, rotazione, appoggio plantare in equinismo e supinazione

16-17 MAGGIO 2025

Intervento chirurgico con Fissatore Esterno Esapodalico

Prima fase

Seconda fase

16-17 MAGGIO 2025

16-17 MAGGIO 2025

**intervento di ortoplastica
lembo vascolarizzato F.E. esapodalico**

ROMA

CONGRESSO NAZIONALE
SOCIETÀ ITALIANA
FISSAZIONE ESTERNA

16-17 MAGGIO 2025

16-17 MAGGIO 2025

16-17 MAGGIO 2025

rimozione F.E.

La fissazione esterna (monolaterale-ibrida-circolare-esapodalica) risulta un valido aiuto nel trattamento delle osteomieliti in eta' pediatrica

L'approccio multidisciplinare e' importante per il risultato finale

ROMA

CONGRESSO NAZIONALE
SOCIETÀ ITALIANA
FISSAZIONE ESTERNA

16-17 MAGGIO 2025

A.S.A.M.I.

Associazione per lo Studio e l'Applicazione del Metodo di Ilizarov

Introduzione alla conoscenza delle metodiche di Ilizarov in ortopedia e traumatologia

A cura di: A. Bianchi-Maiocchi

con la collaborazione di:
G.B. Benedetti - M. Catagni - R. Cattaneo -
L. Tentori - A. Villa

GRAZIE